

Data Alignment, Part II

David N. DeJong
University of Pittsburgh

Spring 2008

Two Issues. #1:

- ▶ Empirical assessments of DSGEs require trend removal.
- ▶ Yet, levels and trends typically convey information regarding model parameterization, through implications regarding steady state relationships.
- ▶ How can we preserve this information, while accomplishing the detrending step?

Issue #2: Upon detrending,

- ▶ Log-linear model representations are in terms of variables expressed as logged deviations from steady state;
- ▶ Non-linear representations are in terms of levels.
- ▶ How can we maintain direct comparability of likelihood functions across alternative representations?

Motivation, cont.

Consider two alternative state-space representations of the optimal growth model:

- ▶ Non-linear
- ▶ Log-Linear

Motivation, cont.

Log-linear:

$$\begin{aligned}x_t &= Fx_{t-1} + e_t, \\x_t &= [\hat{y}, \hat{c}, \hat{i}, \hat{k}, \hat{z}]', \\ \hat{a}_t &= \ln \frac{a_t}{\bar{a}_t}, \\ a_t &= \frac{A_t}{\left(1 + \frac{g}{1-\alpha}\right)^t}.\end{aligned}$$

Observation:

$$X_t = H'x_t + u_t, \quad u_t \sim N(0, \Sigma_u).$$

Motivation, cont.

Non-linear:

$$\left(1 + \frac{g}{1 - \alpha}\right) k'(\tilde{k}_t, \tilde{z}_t) = i(\tilde{k}_t, \tilde{z}_t) + (1 - \delta)k_t$$

$$\log z_t = (1 - \rho) \log(z_0) + \rho \log z_{t-1} + \varepsilon_t$$

$$\tilde{a} = \frac{a_t - \bar{a}}{\omega_a} \in [-1, 1].$$

Observation (if direct comparability is to be maintained across model representations):

$$X_t = H'x_t + u_t, \quad u_t \sim N(0, \Sigma_u).$$

Steady state values:

$$\bar{k} = \left(\frac{\alpha}{(\beta\zeta)^{-1} - (1 - \delta)} \right)^{\frac{1}{1-\alpha}}$$

$$\bar{y} = \bar{k}^\alpha,$$

$$\bar{i} = \left(\delta + \frac{g}{1 - \alpha} \right) \bar{k}$$

$$\bar{c} = \bar{y} - \bar{i}$$

Preserving Information

Let's take $[y, i]$ as observable. A convenient means of representing the model's characterization of these variables is in the form

$$a_t = \bar{a} e^{\left(\frac{g}{1-\alpha}\right)t} e^{\omega_{at}}, \quad a = y, i,$$

where ω_{at} is a deterministic function of the stochastic component ω_t of the TFP process z_t , along with (k_0, z_0) . The mapping from (ω_t, k_0, z_0) to ω_{at} is obtained implicitly from the policy function approximations described above.

Preserving Information, cont.

Recall that model variables are represented in detrended form

$$\frac{a_t}{\left(1 + \frac{g}{1-\alpha}\right)^t} \approx \frac{a_t}{e^{\left(\frac{g}{1-\alpha}\right)t}}.$$

Therefore, in working with the log-linear model approximation,

$$\begin{aligned} \hat{a}_t &= \omega_{at} \\ &= \sigma_k \hat{k}_{t-1} + \sigma_z \hat{z}_{t-1} + e_{ct}; \end{aligned}$$

and in working with the non-linear approximation,

$$\begin{aligned} \frac{a_t}{\left(1 + \frac{g}{1-\alpha}\right)^t} &= \bar{a} e^{\omega_{at}} \\ &= a\left(\tilde{k}_t, \tilde{z}_t\right). \end{aligned}$$

Preserving Information, cont.

Preserving information regarding g .

If the actual data truly followed a stable log-linear trend, detrending could be achieved as part of the model estimation process: for each candidate parameterization, use the relevant (α, g) to construct

$$\frac{a_t}{\left(1 + \frac{g}{1-\alpha}\right)^t}.$$

However, the trend-break behavior we've noted makes this inappropriate.

Instead, we typically work with

$$\ln \frac{a_t}{HP_{at}},$$

which has no trend or intercept term.

Remedy:

Impose $g = (1 - \alpha) \cdot \hat{g}_y$, where \hat{g}_y is the sample estimate of the growth rate of y . (0.00475 for post-war quarterly data).

Preserving Information, cont.

Preserving information regarding steady states.

As noted,

$$\ln \frac{a_t}{HP_{at}}$$

has no trend or intercept term.

Preserving Information, cont.

Remedy:

Restore relative sample values to preserve information regarding steady states:

$$\ln y_t = \ln \frac{Y_t}{HP_{yt}},$$

$$\ln i_t = \ln \frac{I_t}{HP_{it}} + \ln \left(\frac{\bar{i}}{\bar{y}_T} \right),$$

where $\left(\frac{\bar{i}}{\bar{y}_T} \right)$ denotes the sample mean of $\frac{i_t}{y_t}$.

Model Alignment

Assuming interest in maintaining direct comparability of likelihoods across models:

- ▶ For both representations, construct X_t by subtracting logged steady state ratios $\frac{\bar{a}}{\bar{y}}$, $a = y, i$ from logs of the observed data for each proposed model parameterization. The individual elements of X_t are thus

$$\hat{y}_t = \ln \frac{Y_t}{HP_{yt}},$$

$$\hat{i}_t = \ln \frac{I_t}{HP_{it}} + \ln \left(\frac{\bar{i}}{\bar{y}_T} \right) - \ln \frac{\bar{i}}{\bar{y}},$$

- ▶ For the log-linear approximation, X_t is fed directly into the Kalman filter.

Model Alignment, cont.

- ▶ For the non-linear approximation, compare X_t with logged values of the ratio of corresponding model variables to their steady state values:

$$\ln \frac{a(\tilde{k}_t, \tilde{z}_t)}{\bar{a}}, \quad a = y, i.$$

Differences between the elements of X_t and their model counterparts are independent and normally distributed, according to the measurement equation $X_t = H'x_t + u_t$.