

## Chapter 3 – Postulates

1. State of QM system completely specified by wavefunction  $\Psi(x,t)$

$$P(x_0, t_0) = \Psi(x_0, t_0)^* \Psi(x_0, t_0) dx = |\Psi(x_0, t_0)|^2 dx$$

↑  
probability of finding the particle within  $dx$  of  $x_0$  at time  $t_0$

$$\int_{-\infty}^{\infty} |\Psi(x, t)|^2 dx = 1 \quad \longleftarrow \quad \text{probability of finding the particle somewhere}$$

$\Rightarrow \Psi$  is single valued

$\Psi$  and  $\frac{d\Psi}{dx}$  are continuous

$\Psi$  cannot be  $\infty$  over a finite interval

## 2. Each observable is associated with a QM operator

position:  $\hat{x} = x$

momentum:  $\hat{p} = \frac{\hbar}{i} \frac{\partial}{\partial x}$

KE:  $\hat{E}_{kin} = -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} = \frac{\hat{p}^2}{2m}$

PE:  $\hat{E}_{pot} = V(x)$

total E:  $\hat{H} = -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + V(x)$

angular  
momentum:  $\hat{l}_x = \frac{\hbar}{i} \left( y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right)$

....

3. In a single measurement of an observable associated with  $\hat{A}$ , only an eigenvalue of  $\hat{A}$  can be measured.

4. Expectation value: 
$$\langle a \rangle = \frac{\int_{-\infty}^{\infty} \Psi^* \hat{A} \Psi dx}{\int_{-\infty}^{\infty} \Psi^* \Psi dx}$$

Average of the observable  $a$ , if many measurements are done

If  $\Psi$  is an eigenfunction of  $\hat{A}$ , all measurements give the same result

If  $\Psi$  is not an eigenfunction of  $\hat{A}$

$$\Psi = \sum b_n \phi_n(x, t)$$

↑  
eigenfunctions of  $\hat{A}$

$$\langle a \rangle = \sum |b_m|^2 a_m, \quad \text{assuming } \Psi \text{ is normalized}$$

Suppose  $\psi(x) = \frac{1}{2}\phi_1(x) + \frac{\sqrt{3}}{2}\phi_2(x)$ ,  $\phi_1, \phi_2$  being eigenfunctions of  $\hat{A}$

$$\hat{A}\phi_1 = a_1\phi_1, \quad \hat{A}\phi_2 = a_2\phi_2$$

How frequently do we measure  $a_1$ ?  $a_2$ ?

5. The time evolution of a QM system is given by

$$i\hbar \frac{\partial \Psi}{\partial t}(x, t) = \hat{H} \Psi(x, t)$$

If  $\Psi$  is a solution of the time-independent SE

$$\Psi = \psi(x)e^{-iEt/\hbar}$$