



# Selenocysteine: The 21st Amino Acid



Amanda L. Garner

October 5, 2007

# Selenocysteine - Discovery




## Chronology:

- Selenium (Se) initially believed to be toxic (more reactive than sulfur)
- 1954: First report of Se requirement for function of bacterial formate dehydrogenase
- 1959: First paper addressing selenocysteine (Sec)
- 1972: First biochemical studies on role of Se at enzyme level
- 1986: Key study implicating UGA as codon for Sec

Se currently regarded as essential nutrient in humans and mammals (development, immune function, male reproduction, aging process)

# Translational Chain Elongation - Other 20 Amino Acids



# Requirements for Sec Insertion into Protein

1. Sec tRNA<sup>[Ser]<sub>Sec</sub></sup>
2. UGA Codon
3. Selenocysteine Insertion Sequence (SECIS) elements
4. SECIS-binding protein 2 (SBP2)
5. Sec-specific elongation factor (EF<sub>sec</sub>)

# Sec tRNA<sup>[Ser]Sec</sup>

Sec tRNA only known tRNA that governs entire class of proteins *se/C* gene product

Two known structures: 9/4 and 7/5 cloverleaf


90 nt long (longest tRNA known)

1 extra nt in acceptor and T-stems; 2 extra nt in D-stem; longest variable arm

Highly conserved vs. other tRNAs

Occurs as single gene copy in all mammals


Two isoforms: mcm<sup>5</sup>U or mcm<sup>5</sup>Um (final step of Sec tRNA maturation; responsive to Se)



# Sec tRNA<sup>[Ser]Sec</sup>

First aminoacylated with Serine (Ser, S)

Mechanism well known in bacteria but not in eukaryotes



# UGA Codon

UGA = termination codon and Sec codon

Discovered by Böck in 1986 in bacterial formate dehydrogenase (FDH)

*In vivo* studies confirmed that Sec tRNA<sup>[Ser]<sup>Sec</sup> recognizes the UGA codon in both bacteria and eukaryotes (definitive that Sec is 21<sup>st</sup> amino acid)</sup>

What determines whether UGA will code for termination or Sec?

# SECIS Elements


Bacteria - found as 40 base  
sequence downstream from UGA

Eukaryotes - found in 3'-UTR

Primary sequence not conserved  
(except bases in red)

Main functional site is non-Watson-  
Crick quartet

Responsible for Sec insertion if can  
access ribosome




# SECIS Binding Protein 2 (SBP2)

Forms strong complex with SECIS quartet and sequence preceding (not apical loop)

Stably associated with 28S rRNA of ribosome (preselects for Sec insertion)

Also binds E<sub>f</sub>Sec to recruit Sec tRNA<sup>[Ser]<sub>Sec</sub></sup>

mcm<sup>5</sup>Um may be involved in initiating complexation




# Elongation Factor EFsec

se/B gene product

Recognizes extra bp in acceptor stem of Sec tRNA<sup>[Ser]Sec</sup>

Ready to insert Sec into protein in response to UGA codon



# Selenocysteine - Evolution

Evolution - believed to be a latter addition to the existing genetic code of 20 amino acids

Sec found in bacteria, archaea, and eukaryotes

Bacteria/archaea - found in proteins responsible for catabolic processes

Eukaryotes - found in proteins responsible for anabolic and regulatory pathways

Increased use of Sec in proteins of higher vertebrates

\*Sec is an evolutionary advancement

# Examples of Known Selenoproteins

## Prokaryotes:

1. Glycine reductase (first known selenoprotein; 1973)
2. Formate dehydrogenase
3. Selenophosphate synthetase


## Eukaryotes:

1. Glutathione peroxidase (first known selenoprotein; 1973)
2. Thioredoxin reductase
3. Thyroid hormone deiodinase
4. Selenophosphate synthetase
5. Selenoprotein P

Only 12 eukaryote selenoproteins with known function  
25 total selenoproteins contained in the human genome

# Glutathione Peroxidase

Key enzyme in oxidative stress by scavenging peroxides



# Glutathione Peroxidase Structure

QuickTime™ and a  
TIFF (Uncompressed) decompressor  
are needed to see this picture.

From human plasma

RCSB Protein Data Bank

# Thioredoxin/Thioredoxin Reductase

## Thioredoxin (Trx)

Major cellular protein disulfide reductase

Numerous important roles in mammals:

1. DNA synthesis (binds T7 DNA polymerase)
2. Redox regulation of transcription factors
3. Regulation of apoptosis (bind apoptosis signaling kinase)
4. Immunomodulation
5. Pregnancy and Birth
6. CNS

Contains important Trx fold (5  $\beta$ -strands and 4  $\alpha$ -helices) and Cys-Gly-Pro-Cys active site sequence


# Thioredoxin Structure

QuickTime™ and a  
TIFF (Uncompressed) decompressor  
are needed to see this picture.

# Thioredoxin/Thioredoxin Reductase

Thioredoxin reductase (TrxR; 1996, Stadtman)

Se essential for activity; Gly-Cys-Sec-Gly motif conserved



Redox potential of  
each couple unknown

# Thioredoxin Reductase Structure

QuickTime™ and a  
TIFF (Uncompressed) decompressor  
are needed to see this picture.

# Selenoprotein P

Abundant extracellular protein rich in Sec

Identified in 1977 in rat plasma; 1982 shown to contain Sec

Only expressed in vertebrates

First known protein with >1 Sec (10 total in human)

Two Sec-containing domains:

1. N-terminal domain - UxxC redox motif


2. C-terminal domain - 9 U and 10 C (no redox motif)

3D structure unknown but contains 3 Se-S motifs and 2 S-S motifs; 2 SECIS elements

Exhibits weak GPx activity but function unknown

Speculated as Se transport protein but no redox activity

# Organoselenium Pharmacology



Ebselen: GPx mimetic agent (anti-oxidant), anti-inflammatory activity, neuroprotection

Ph<sub>2</sub>Se<sub>2</sub>: GPx mimic, anti-inflammatory activity, antinociceptive activity (greater than Ebselen)

Selenocyanate: Chemopreventative (colon, breast, lung, liver, intestine, oral tissue carcinomas)

# Uniqueness of Selenium

Cys and Sec least abundant aa in proteins

In adult human, 140 g S but only mg quantities of Se

Se more polarizable (better nucleophile, leaving group and lower redox potential)

RSeH: pKa = 5.7 vs. RSH: pKa = 8.5 (~1 pKa unit higher in protein)

RSeH largely deprotonated at physiological pH

# Redox Properties of RSeH/R<sub>2</sub>Se<sub>2</sub>



Only S nucleophile known to reduce Se-Se is dithiothreitol (DTT, -323 mV)

GSH and cysteine also shown to reduce but at concentrations 10<sup>3</sup> greater than [Se-Se]

Trx synthesized as double U mutant forming Se-Se

- Redox properties substantially different from native protein (could not be reduced by β-mercaptoethanol)

# Cys→Sec Mutants

Mutation typically shows catalytic enhancement


Examples:

1. Formate dehydrogenase:  $k_{\text{cat}} = 9 \text{ s}^{-1}$  (C);  $2800 \text{ s}^{-1}$  (U)
2. Deiodinase: C only 10-20% as active as U
3. GPx: C has insignificant activity

Although more active, little known about kinetic role of Se in enzyme active site

Implications for Sec as evolutionary enhancement

# Synthetic Seleno-Glutaredoxin3 Analogues



Redox potentials:

$^{11}\text{Cxx}^{14}\text{C}$ : -194 mV

C11U: -260 mV

C14U: -275 mV

$^{11}\text{Uxx}^{14}\text{U}$ : -309 mV

Sec more active  
than Cys

(Reference: Trx = -270 mV; GSH = -240 mV)

~9000-fold enhancement for double mutant (relative to  $k_{-1}$ )

Equilibrium studies show ~19% Se-Se reduced suggesting Se-Se could have role in biological catalysis if found in natural protein

# Identification and Characterization of a Selenoprotein Family Containing a Diselenide Bond

Identified SelL, only known selenoprotein containing 2 Sec in UxxU redox motif

Only present in aquatic organisms:

Eukaryotes - 11 fish, 2 ascidian, 2 crustacean, 1 mollusk

Prokaryotes - 2 bacteria

Unknown why only found in aquatic organisms

Close Cys homolog found in mammals

# Identification and Characterization of a Selenoprotein Family Containing a Diselenide Bond

## Structure:

Prx-like domain in C-terminus

Conserved UxxU in N-terminus

2 Sec in loop between  $\beta$ -strand and  $\alpha$ -helix (Trx-like)

Single SECIS element for 2 Sec (contains CA vs. AA  
as unpaired nt in apical loop)

SelL Structure: Figure 1A, 1B (2 Sec shown in red)

# Identification and Characterization of a Selenoprotein Family Containing a Diselenide Bond

1. Determine if single SECIS element can direct addition of 2 Sec

Express 2 prokaryotic Sell sequences in *E. coli* via GST-fused fragment containing both UGA codons, SECIS element, and C-terminal His tag  
(Figure 2A, 2B, 2E)

# Identification and Characterization of a Selenoprotein Family Containing a Diselenide Bond

## 2. Confirmation of 2 Sec by mass spectrometry

Method = LC-ESI MS/MS

Mass (from neutral, +2, and +3) = 1039.3 with correct isotopic distribution

(Figure 3A)

# Identification and Characterization of a Selenoprotein Family Containing a Diselenide Bond

3. Expression in eukaryote, *Danio rerio* (zebrafish)  
Constructs with C-terminal His-tagged SelL (and mutants ULPC, CLPU, CLPC)  
Transfected into HEK293 and NIH 3T3 cells  
(Figure 2C, 3C)

# Identification and Characterization of a Selenoprotein Family Containing a Diselenide Bond

## 4. Subcellular localization

Expressed GST-fused *D. rerio* SelL in NIH 3T3 cells  
Cys mutant with GFP at terminus

Observe cytosolic localization

(Supporting Information)

# Identification and Characterization of a Selenoprotein Family Containing a Diselenide Bond

## 5. Redox properties of Sell

Expressed in HEK293 cells (Cys→Ser)

a. *Method 1* = reduced all S-S with DTT and alkylated; analyze by SDS-PAGE

Result: all other selenoproteins but mutated Sell shifted (wt Sell did shift)

b. *Method 2* = pretreat with *N*-ethylmaleimide, then DTT and alkylation

Result: no Sell shift observed (Se-Se bond!)

(Supporting Information)

# Identification and Characterization of a Selenoprotein Family Containing a Diselenide Bond

## Summary

Isolated first Se-Se containing natural protein, Sell  
Demonstrated a single SECIS element can direct the  
incorporation of 2 Sec

Many questions remain:

Does a catalytic role exist for Se-Se in proteins?

Why is Sell only found in aquatic organisms?

Do other proteins contain this interesting motif?