



Mercury, Arsenic and Selenium in White Bass fillet caught in the Allegheny and Monongahela Rivers near Pittsburgh PA; Comparisons with store-bought fish from Canadian Lake Erie

APHA Conference 2007-Paper #161184

**Conrad Dan Volz, DrPH, MPH¹, Nancy
Sussman, PhD¹, Devra Lee Davis, PhD, MPH²,
Maryann Donovan, MPH, PhD², Yan Liu, BS
Env Eng, MPH¹, Sean Brady, BS, MA³, Karen
Gainey⁴, and Jeanne Zborowski, PhD, MS⁵**

**(1) Department of Environmental and Occupational
Health, Graduate School of Public Health,
University of Pittsburgh (2) Center for
Environmental Oncology, University of Pittsburgh
Cancer Institute (3) Venture Outdoors (4) Kamsport
(5) Department of Epidemiology, Graduate School
of Public Health, University of Pittsburgh**

Why should we be concerned about toxic metals, metalloids and elements in White

Bass (*Morone chrysops*)

- The white bass is a freshwater, temperate member of the bass family Moronidae. They range throughout the central USA, west of the Appalachians into the Great Lakes and in the riverine systems in the Ohio and Mississippi river valleys.
- White bass are a favorite sports fishing species and schools of white bass feeding on shad generate considerable excitement amongst both semi-subsistence and recreational anglers.
- River-caught white bass are excellent table fare. Since an angler can literally catch a hundred at a time if spooning within a school, large quantities are eaten at fish fries and are stored in the freezer for future consumption.
- There now exist commercial fisheries for white bass in Canadian Lake Erie. These fish are sold in fish markets in Pittsburgh PA, other cities within the USA and in Canada.

Initial Hypothesis

- **Mercury (Hg), arsenic (As), and selenium (Se) levels in white bass fillet (in mg/kg wet weight) will be higher from specimens taken near legacy Iron and Steel pollution sites on the Monongahela River than from the less industrially impacted Allegheny River site, even though both are in the same river pool system.**
- **Hg, As and Se levels in white bass fillet will be higher in river-caught samples from the Pittsburgh Pool, due to legacy pollution and nearby power plants and municipal pollution sources, than white bass commercially sold in the market and branded as Canadian Lake Erie in origin.**



Objective A:


To determine if white bass fillet concentrations of mercury, arsenic and/or selenium vary by locations of catch, within the Pittsburgh Pool.



Objective B:

To determine if mercury, arsenic and/or selenium concentrations in white bass fillet vary between river caught and store-bought fish (Canadian Lake Erie).

Locations of White Bass Catch Near Pittsburgh and Store Bought in Lake Erie



Three Rivers Area

Highland Park

Approximate range of Store Bought-Lake Erie fish, relative to Pittsburgh, Pennsylvania

Monongahela River near the U. S. Steel Edger Thompson Works

Volz et al.,2007; APHA Conference: <http://www.pitt.edu/~cdv5/>

Methods

White bass were caught by rod and reel method using a Community Based Participatory Research approach. Local anglers were recruited to catch fish. White bass could not be caught at Point State park in the Pittsburgh Pool and at Kittanning Dam on the Allegheny, presumably because these habitats are not conducive to permanent resident fish populations.

The geo-coordinates of all fish collection locations were determined on a Garmin, GPS Map 60CSx device and were used in ArcView 9.2 map production.

10-35 white bass were caught at each location and 10 white bass were store-bought-coming from Canadian Lake Erie.

0.5 gram of white bass fillet was microwave digested in nitric acid. As and Se were analyzed by collision cell ICP-MS with calibration by the method of standard additions. Mercury was analyzed by isotope dilution cold vapor ICP-MS.

Data were log-transformed and analyzed by ANOVA with Tukey post hoc tests using SPSS 15.0.




Descriptive Statistics, Mercury (Hg) Concentrations in Fish Fillet by Catch Location and in Store Bought Fish

Catch Location	n	Mean Concentration mg/kg* 95% CI of the Mean	Median Concentration mg/kg Range
Pittsburgh Pool- Monongahela River at Braddock Dam	1 0	0.041 0.025-0.067	0.033 0.026-0.081
Pittsburgh Pool- Allegheny River at Highland Park Dam	3 5	0.034 0.031-0.037	0.033 0.021-0.070
Store Bought- Caught in Canadian Lake Erie	1 0	0.137 0.126-0.148	0.134 0.118-0.163

*mg/kg is equivalent to ppm, all samples are by wet weight.

Volz, et al. 2007; APHA Conference: <http://www.pitt.edu/~cdv5/>

95% Confidence Interval of the Mean Mercury Concentration (ppm) in White Bass Fillet from the Pittsburgh Pool vs Store-bought (Canadian Lake Erie)



Pittsburgh Pool vs Other

Descriptive Statistics, Arsenic (As) Concentrations in Fish Fillet by Catch Location and in Store Bought Fish


Catch Location	n	Mean As Concentration mg/kg* 95% CI of the Mean	Median As Concentration mg/kg Range
Pittsburgh Pool- Monongahela River at Braddock Dam	10	0.111 0.079 - 0.156	0.112 0.049 - 0.194
Pittsburgh Pool- Allegheny River at Highland Park Dam	35	0.126 0.111 - .143	0.119 0.071 - .359
Store Bought- Caught in Canadian Lake Erie	10	0.206 .165 -.257	0.202 .127 - .361

*mg/kg is equivalent to ppm, all samples are by wet weight.

Volz, et al. 2007; APHA Conference: <http://www.pitt.edu/~cdv5/>



95% Confidence Interval of the Mean Arsenic Concentration (ppm) in White Bass Fillet from the Pittsburgh Pool vs Store-bought (Canadian Lake Erie)



Pittsburgh Pool vs Other

Descriptive Statistics, Selenium (Se) Concentrations in Fish Fillet by Catch Location and in Store Bought Fish

Catch Location	n	Mean Concentration Se mg/kg* 95% CI of the Mean	Median Se Concentration mg/kg Range
Pittsburgh Pool- Monongahela River at Braddock Dam	10	0.547 0.444 - 0.674	0.574 0.334 - 0.780
Pittsburgh Pool- Allegheny River at Highland Park Dam	35	0.570 0.508 - 0.640	0.541 0.376 - 0.739
Store Bought- Caught in Canadian Lake Erie	10	1.01 0.865-1.177	1.11 0.670 - 1.20

*mg/kg is equivalent to ppm, all samples are by wet weight.

White Bass, ANOVA Tukey Post Hoc Comparisons



**White Bass
Fillet Mercury
Concentration**

Lower

Pittsburgh Pool-Hg

Median: 0.033 ppm

Kruskal Wallis

$\chi^2 = 22.2$

$p = .0001$

ANOVA

$p = .0001$

95%CI of median ratio

2.98 to 5.45

High

**Local Store Bought White Bass-Hg
Canadian Lake Erie**

Median: 0.134 ppm

**White Bass
Fillet Arsenic
Concentration**

Lower



High

Pittsburgh Pool-As

Median: 0.119 ppm

Kruskal Wallis

$\chi^2 = 13.4$

$p = .001$



ANOVA

$p = .002$

95%CI of median ratio

1.24 to 2.76

**Local Store Bought White Bass-As
Wild Caught-Canadian Lake Erie**

Median: 0.202 ppm

**White Bass
Fillet Selenium
Concentration**

Lower

Pittsburgh Pool-Se
Median: 0.57 ppm

Kruskal Wallis

$\chi^2 = 19.3$

$p = .0001$



ANOVA

$p = .0001$

95%CI of median ratio

1.32 to 2.58

High

**Local Store Bought White Bass-Se
Wild Caught-Canadian Lake Erie**
Median: 1.11 ppm



Conclusions

- **We reject our hypothesis that there is a significant difference in Hg, As and/or Se levels in fillet tissue of white bass from legacy contaminated areas of the Monongahela River compared to the less industrially impacted Allegheny River site, within the Pittsburgh Pool.**
- ✓ **The Pittsburgh Pool may be considered a locked water system due to the lock and dam system, which essentially makes a lake around Pittsburgh including portions of the Allegheny, Monongahela and Ohio Rivers.**
- ✓ **Pollution discharged into the Pittsburgh Pool from industrialized sources and sewer overflows may have become somewhat evenly distributed throughout the water body-especially over time.**
- ✓ **Fish caught at each location within the Pittsburgh Pool may range throughout the pool. This is especially true for white bass which travel in large schools and congregate near dams where the dissolved oxygen level is maximized.**



Conclusions

- **We reject our hypothesis that river-caught fish have less Hg, As and/or Se than commercially sold market fish. Store-bought fish had 4.1, 1.7, and 1.9 times the median Hg, As, and Se levels found in river-caught fish, respectively.**
- ✓ **The store-bought fish from Canadian Lake Erie were significantly larger than the river-caught fish, which helps explain why there is more bioconcentrated Hg and Se in Lake Erie caught fish.**
- ✓ **The ecological habitat and food chain for riverine and Lake Erie (big lake) white bass is considerably different. There is a more complex food chain in the Great Lakes compared to the Upper Ohio River drainages, this includes more trophic levels in the lake system and with them- more opportunity for bioaccumulation and contaminant magnification.**



Public Health Implications



- **Location, location, location-just as in real estate the location of fish catch can help us to make informed hypothesis regarding the contaminants-toxic metals and metalloids, elements and even persistent organic pollutants that may be in fish tissue.**
- **Lake Erie has received immense legacy pollution assaults from our industrial cities including Detroit, Toledo, Cleveland, Buffalo and Windsor, Canada. There is ongoing deposition of power plant air and water emissions directly into Lake Erie and into its watershed.**

Acknowledgements

- **The project is funded by the Highmark Foundation, Healthy People-Healthy Communities Project, the DSF Charitable Trust and the Heinz Endowments through the Center for Environmental Oncology of the University of Pittsburgh Cancer Institute.**
- **CDV, DLD and JZ receive funding through the Centers for Disease Control and Prevention's, Environmental Public Health Tracking (EPHT) Network through the University of Pittsburgh's Academic Center for Excellence in EPHT.**
- **Program partners included Venture Outdoors, Clean Water Action, and the Bassmasters Groups of Western Pennsylvania.**
- **Special thanks to Mr. Paul Caruso, Ms. Karen Gainey, Mr. Matt Weaver and many individual anglers in the Southwestern PA area.**

Conrad (Dan) Volz, DrPH, MPH, Bridgeside Point, 100 Technology Drive, Suite 564, BRIDG, Pittsburgh, PA 15219-3130. office 412-648-8541 cdv5@pitt.edu

Volz et al.,2007; APHA Conference: <http://www.pitt.edu/~cdv5/>

